

# Multi-Quarks and Two-Baryon Interaction in Lattice QCD

F. Okiharu\*, H. Suganuma<sup>†</sup>, T. T. Takahashi\*\* and T. Doi<sup>‡</sup>

\* *Nihon University, 1-8-14 Kanda Surugadai, Chiyoda, Tokyo 101-8308, Japan*

<sup>†</sup> *Dept. of Phys., Kyoto University, Kitashirakawa, Sakyo, Kyoto 606-8502, Japan*

\*\* *YITP, Kyoto University, Kitashirakawa, Sakyo, Kyoto 606-8502, Japan*

<sup>‡</sup> *RIKEN-BNL Research Center, BNL, Upton, New York 11973, USA*

**Abstract.** We study multi-quark (3Q,4Q,5Q) systems in lattice QCD. We perform the detailed studies of multi-quark potentials in lattice QCD to clarify the inter-quark interaction in multi-quark systems. We find that all the multi-quark potentials are well described by the OGE Coulomb plus multi-Y-type linear potential, i.e., the multi-Y Ansatz. For multi-quark systems, we observe lattice QCD evidences of “flip-flop”, i.e., flux-tube recombination. These lattice QCD studies give an important bridge between elementary particle physics and nuclear physics.


## INTRODUCTION

The inter-quark force [1, 2, 3] is one of the most important quantities in hadron physics to understand hadron properties at the quark-gluon level, and also plays an essential role to reveal properties of multi-quarks [4, 5, 6]. Nevertheless, before our studies, nobody knew the exact form of the confinement force in the multi-quark systems directly from QCD. In fact, some hypothetical forms of the inter-quark potential have been used in all the quark model calculations. Then, the lattice QCD study of the inter-quark interaction is quite desired, and it leads to the proper Hamiltonian and a guideline to construct the QCD-based quark model for multi-quark systems.

We show in Fig.1 our global strategy to understand hadron properties from QCD. One way is the direct lattice QCD calculations for low-lying hadron masses [7] and simple hadron matrix elements, although the wave function cannot be obtained in the path-integral formalism and the practically calculable quantities are severely limited. The other way is to construct the quark model from QCD. From the analysis of the inter-quark forces in lattice QCD, we extract the quark-model Hamiltonian. Through the quark model calculation, one can obtain the quark wave-function of hadrons and more complicated properties of hadrons.

## MULTI-QUARK INTERACTION FROM LATTICE QCD

In order to clarify the inter-quark force in multi-quark systems, we study the static multi-quark potentials systematically in lattice QCD using the multi-quark Wilson loops [1, 2].




**FIGURE 1.** Our global strategy to understand the hadron properties from QCD.

As the results, we clarify that the multi-quark potential is well described as

$$V = \frac{g^2}{4\pi} \sum_{i < j} \frac{T_i^a T_j^b}{|\mathbf{r}_i - \mathbf{r}_j|} + \sigma L_{\min} + C, \quad (1)$$

where  $L_{\min}$  is the minimal value of the total length of flux-tubes linking the static quarks.

From lattice QCD, the QQ potential  $V_{Q\bar{Q}}$  is known to be well described by this form [1, 3]. For the 3Q potential  $V_{3Q}$ , our accurate calculations clarify that  $V_{3Q}$  is well described by the OGE Coulomb plus Y-type linear potential, i.e., the Y-Ansatz[1, 2], which supports the Y-shaped flux-tube formation [1, 2, 8]. Note that the Y-type linear potential implies existence of the three-body interaction. (In usual many-body systems, the main interaction is described by two-body interactions and the three-body interaction is a higher-order contribution. In contrast, as is clarified by lattice QCD study, the quark confinement force in baryons is a genuine three-body interaction.)



**FIGURE 2.** The lattice QCD results for the 3Q, 4Q and 5Q potentials. (a) the 3Q confinement potential (the Coulomb-subtracted 3Q potential)  $V_{3Q}^{\text{conf}}$  v.s.  $L_{\min}^{3Q}$ , (b) “flip-flop” in  $V_{4Q}$ , which takes the smaller energy of the multi-Y 4Q state or the two-meson state, (c)  $V_{5Q}$  v.s.  $L_{\min}^{5Q}$ . (For details, see Ref.[2].)

We study the 4Q potential  $V_{4Q}$  in lattice QCD for QQ- $\bar{Q}\bar{Q}$  configurations [2]. When QQ and  $\bar{Q}\bar{Q}$  are well separated,  $V_{4Q}$  is well described by the OGE Coulomb plus multi-Y Ansatz with a single connected flux-tube. When the nearest QQ pair is spatially close,  $V_{4Q}$  is well described by the sum of two QQ potentials, which indicates a “two-meson” state. Thus,  $V_{4Q}$  is found to take the smaller energy of two states. In other words, we observe a clear lattice QCD evidence of “flip-flop”, i.e., flux-tube recombination between the two states. As for 5Q systems, we study QQ-Q-QQ systems and find that the 5Q potential  $V_{5Q}$  is well described by the OGE Coulomb plus multi-Y Ansatz[2].

Thus, from a series of our lattice QCD studies, the inter-quark potential is clarified to consist of the OGE Coulomb part and the flux-tube-type linear confinement part in both mesons, baryons and multi-quark hadrons [1, 2].



**FIGURE 3.** The flux-tube formation in multi-quarks indicated by our lattice QCD results.

From the comparison among the  $Q\bar{Q}$ , 3Q, 4Q and 5Q potentials in lattice QCD, we find the universality of the string tension  $\sigma$  [1, 2] as  $\sigma_{Q\bar{Q}} \simeq \sigma_{3Q} \simeq \sigma_{4Q} \simeq \sigma_{5Q}$ , and the OGE Coulomb coefficient, i.e.,  $g \simeq 1.6$ . Here, the OGE Coulomb term is considered to originate from the OGE process, which plays the dominant role at short distances, where perturbative QCD is applicable. The flux-tube-type linear confinement physically indicates the flux-tube picture, where quarks and antiquarks are linked by the one-dimensional squeezed color-electric flux-tube with the string tension  $\sigma \simeq 0.89\text{GeV/fm}$ .

To conclude, the inter-quark interaction would be generally described by the sum of the OGE part and the flux-tube-type linear confinement part with the universal string tension  $\sigma$ . Thus, based on the lattice QCD results, we propose the proper quark-model Hamiltonian  $\hat{H}$  for multi-quark hadrons as

$$\hat{H} = \sum_i \sqrt{\hat{\mathbf{p}}_i^2 + M_i^2} + \sum_{i<j} V_{\text{OGE}}(\mathbf{r}_i - \mathbf{r}_j) + \sigma L_{\text{min}}, \quad (2)$$

where  $V_{\text{OGE}}$  denotes the OGE potential and  $M_i$  the constituent quark mass.

To summarize, we have performed the multi-quark potential in lattice QCD to clarify the inter-quark interaction in multi-quark systems. We have found that the 3Q potential is well described by the Y-Ansatz. For 4Q and 5Q systems,  $V_{4Q}$  and  $V_{5Q}$  are well described by the OGE Coulomb plus multi-Y Ansatz, which indicates the flux-tube picture even for multi-quarks. We have observed a clear lattice QCD evidence of “flip-flop”.

As a successive work, we are now investigating the nuclear force in lattice QCD [9].

The lattice calculations have been done on NEC-SX5(Osaka U.) and SR8000(KEK).

## REFERENCES

1. T.T. Takahashi, H. Suganuma et al., Phys. Rev. Lett. **86** (2001) 18; Phys. Rev. **D65** (2002) 114509; Phys. Rev. Lett. **90** (2003) 182001; Phys. Rev. **D70** (2004) 074506.
2. F. Okiharu, H. Suganuma et al., Phys. Rev. **D72** (2005) 014505; Phys. Rev. Lett. **94** (2005) 192001.
3. H. J. Rothe, “Lattice Gauge Theories”, 3rd edition (World Scientific, 2005).
4. Belle Collab., Phys. Rev. Lett. **91** (2003) 262001; *ibid.* **91** (2003) 262002; *ibid.* **94** (2005) 182002.
5. BaBar Collaboration, Phys. Rev. Lett. **90** (2003) 242001; *ibid.* **93** (2004) 041801.
6. For recent experimental status of pentaquarks, e.g., K. Hicks, Prog. Part. Nucl. Phys. **55** (2005) 647.
7. N. Ishii et al., Phys. Rev. **D71** (2005) 034001; Phys. Rev. **D72** (2005) 074503.
8. H. Ichie, V. Bornyakov, T. Streuer and G. Schierholz, Nucl. Phys. **A721** (2003) 899.
9. T. Doi et al., this proceedings; T.T. Takahashi et al., this proceedings.